

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--

Pearson Edexcel Level 3 GCE

Friday 19 May 2023

Afternoon

Paper
reference

8FM0/26

Further Mathematics
Advanced Subsidiary
Further Mathematics options
26: Further Mechanics 2
(Part of option J)

You must have:

Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Unless otherwise indicated, whenever a value of g is required, take $g = 9.8 \text{ m s}^{-2}$ and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- The total mark for this part of the examination is 40. There are 4 questions.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P72812A

©2023 Pearson Education Ltd.

N:1/1/1/



Pearson

1. Three particles of masses $4m$, $2m$ and km are placed at the points with coordinates $(-3, -1)$, $(6, 1)$ and $(-1, 5)$ respectively.

Given that the centre of mass of the three particles is at the point with coordinates (\bar{x}, \bar{y})

(a) show that $\bar{x} = \frac{-k}{k+6}$ (3)

(b) find \bar{y} in terms of k . (2)

Given that the centre of mass of the three particles lies on the line with equation $y = 2x + 3$

(c) find the value of k . (2)

A fourth particle is placed at the point with coordinates $(\lambda, 4)$.

Given that the centre of mass of the four particles also lies on the line with equation $y = 2x + 3$

(d) find the value of λ . (2)



Question 1 continued

Lined writing area for the answer to Question 1.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 1 continued

Lined writing area for the answer to Question 1.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



2. A particle P is moving along the x -axis.
At time t seconds, $t \geq 0$, P has acceleration $a \text{ m s}^{-2}$ and velocity $v \text{ m s}^{-1}$ in the direction of x increasing, where

$$v = e^{2t} + 6e^t - kt$$

and k is a positive constant.

When $t = \ln 2$, $a = 0$

- (a) Find the value of k .

(4)

When $t = 0$, the particle passes through the fixed point A .

When $t = \ln 2$, the particle is d metres from A .

- (b) Showing all stages of your working, find the value of d correct to 2 significant figures.

[Solutions relying entirely on calculator technology are not acceptable.]

(4)



Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Ruled lines for writing.

(Total for Question 2 is 8 marks)



3. A girl is cycling round a circular track.
The girl and her bicycle have a combined mass of 55 kg.
The coefficient of friction between the track surface and the tyres of the bicycle is μ .

The track is banked at an angle of 15° to the horizontal.

The girl and her bicycle are modelled as a particle moving in a horizontal circle of radius 50 m

The minimum speed at which the girl can cycle round this circle without slipping is 4.5 m s^{-1}

Using the model, find the value of μ .

(9)



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Handwriting practice lines for Question 3 continued.



P 7 2 8 1 2 A 0 9 1 6



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined area for writing the answer to Question 3.



4.




Figure 1

A uniform triangular lamina ABC is isosceles, with $AC = BC$. The midpoint of AB is M . The length of AB is $18a$ and the length of CM is $18a$.

The triangular lamina CDE , with $DE = 6a$ and $CD = 12a$, has ED parallel to AB and MDC is a straight line.

Triangle CDE is removed from triangle ABC to form the lamina L , shown shaded in Figure 1.

The distance of the centre of mass of L from MC is d .

- (a) Show that $d = \frac{4}{7}a$ (4)

The lamina L is suspended by two light inextensible strings. One string is attached to L at A and the other string is attached to L at B .

The lamina hangs in equilibrium in a vertical plane with the strings vertical and AB horizontal.

The weight of L is W

- (b) Find, in terms of W , the tension in the string attached to L at B (3)

The string attached to L at B breaks, so that L is now suspended from A .

When L is hanging in equilibrium in a vertical plane, the angle between AB and the downward vertical through A is θ°

- (c) Find the value of θ (7)



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined writing area for the response to Question 4.



P 7 2 8 1 2 A 0 1 5 1 6

