

CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International Advanced Subsidiary and Advanced Level

MARK SCHEME for the May/June 2015 series

9709 MATHEMATICS

9709/11

Paper 1, maximum raw mark 75

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2015 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2015	9709	11

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol \checkmark implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2015	9709	11

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
CWO	Correct Working Only - often written by a “fortuitous” answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR–1	A penalty of MR–1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through” marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR–2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA–1	This is deducted from A or B marks in the case of premature approximation. The PA–1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2015	9709	11

1	<p>θ is obtuse, $\sin \theta = k$</p> <p>(i) $\cos \theta = -\sqrt{1 - k^2}$</p> <p>(ii) $\tan \theta = \frac{\sin \theta}{\cos \theta}$ used $\rightarrow \tan \theta = -\frac{k}{\sqrt{1 - k^2}}$ aef</p> <p>(iii) $\sin(\theta + \pi) = -k$</p>	<p>B1 [1]</p> <p>M1</p> <p>A1[✓] [2]</p> <p>B1 [1]</p>	<p>cao</p> <p>Used, attempt at cosine seen in (i)</p> <p>Ft for their cosine as a function of k only, from part (i)</p> <p>cao</p>
2	<p>$y = 2x^2$, $X(-2, 0)$ and $P(p, 0)$</p> <p>(i) $A = \frac{1}{2} \times (2 + p) \times 2p^2 (= 2p^2 + p^3)$</p> <p>(ii) $\frac{dA}{dp} = 4p + 3p^2$ $\frac{dA}{dt} = \frac{dA}{dp} \times \frac{dp}{dt} = 0.02 \times 20 = 0.4$ or $\frac{dA}{dt} = 4p \frac{dp}{dt} + 3p^2 \frac{dp}{dt}$</p>	<p>M1 A1 [2]</p> <p>B1</p> <p>M1 A1 [3]</p>	<p>Attempt at base and height in terms of p and use of $\frac{bh}{2}$</p> <p>cao</p> <p>any correct method, cao</p>
3	<p>$(1 - x)^2(1 + 2x)^6$.</p> <p>(i) (a) $(1 - x)^6 = 1 - 6x + 15x^2$</p> <p>(b) $(1 + 2x)^6 = 1 + 12x + 60x^2$</p> <p>(ii) Product of (a) and (b) with >1 term $\rightarrow 60 - 72 + 15 = 3$</p>	<p>B2,1 [2]</p> <p>B2,1 [2]</p> <p>M1 DM1A1 [3]</p>	<p>-1 each error</p> <p>-1 each error SC B1 only, in each part, for all 3 correct descending powers SC only one penalty for omission of the '1' in each expansion</p> <p>Must be 2 or more products M1 exactly 3 products. cao, condone $3x^2$</p>

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2015	9709	11

<p>4</p> <p>(i)</p>	$\overrightarrow{OA} = \begin{pmatrix} 3 \\ 0 \\ -4 \end{pmatrix}, \overrightarrow{OB} = \begin{pmatrix} 6 \\ -3 \\ 2 \end{pmatrix}, \overrightarrow{OC} = \begin{pmatrix} k \\ -2k \\ 2k-3 \end{pmatrix}$ <p>$OA \cdot OB = 18 - 8 = 10$ Modulus of $OA = 5$, of $OB = 7$ Angle $AOB = \cos^{-1}\left(\frac{10}{35}\right)$ aef $\rightarrow \frac{10}{35}$ or $\frac{2}{7}$</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>[3]</p>	<p>Use of $x_1x_2 + y_1y_2 + z_1z_2$</p> <p>All linked with modulus cao, (if angle given, no penalty), correct angle implies correct cosine</p>
<p>(ii)</p>	$\overrightarrow{AB} = \mathbf{b} - \mathbf{a} = \begin{pmatrix} 3 \\ -3 \\ 6 \end{pmatrix}$ <p>$k^2 + 4k^2 + (2k - 3)^2 = 9 + 9 + 36$ $\rightarrow 9k^2 - 12k - 45 (= 0)$ $\rightarrow k = 3$ or $k = -\frac{5}{3}$</p>	<p>B1</p> <p>M1</p> <p>DM1</p> <p>A1</p> <p>[4]</p>	<p>allow for $\mathbf{a} - \mathbf{b}$</p> <p>Correct use of moduli using their AB obtains 3 term quadratic. cao</p>
<p>5</p> <p>(i)</p> <p>(ii)</p> <p>(iii)</p>	<p>$24 = r + r + r\theta$ $\rightarrow \theta = \frac{24 - 2r}{r}$ $A = \frac{1}{2} r^2 \theta = \frac{24r}{2} - r^2 = 12r - r^2$. aef, ag</p> <p>$(A =) 36 - (r - 6)^2$</p> <p>Greatest value of $A = 36$ $(r = 6) \rightarrow \theta = 2$</p>	<p>M1</p> <p>M1A1</p> <p>[3]</p> <p>B1 B1</p> <p>[2]</p> <p>B1^{ft}</p> <p>B1</p> <p>[2]</p>	<p>(May not use θ)</p> <p>Attempt at $s = r\theta$ linked with 24 and r</p> <p>Uses A formula with θ as $f(r)$. cao</p> <p>cao</p> <p>Ft on (ii).</p> <p>cao, may use calculus or the discriminant on $12r - r^2$</p>

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2015	9709	11

<p>6 (i)</p> <p>$y - 2t = -2(x - 3t)(y + 2x = 8t)$ Set x to 0 $\rightarrow B(0, 8t)$ Set y to 0 $\rightarrow A(4t, 0)$ $\rightarrow \text{Area} = 16t^2$</p> <p>(ii)</p> <p>$m = \frac{1}{2}$ $\rightarrow y - 2t = \frac{1}{2}(x - 3t)(2y = x + t)$ Set y to 0 $\rightarrow C(-t, 0)$ Midpoint of CP is (t, t) This lies on the line $y = x$.</p>	<p>M1</p> <p>M1 A1 [3]</p> <p>B1</p> <p>M1 A1</p> <p>A1 [4]</p>	<p>Unsimplified or equivalent forms</p> <p>Attempt at both A and B, then using cao</p> <p>cao</p> <p>Unsimplified or equivalent forms co</p> <p>correctly shown.</p>
<p>7 (a)</p> <p>$ar^2 = \frac{1}{3}, ar^3 = \frac{2}{9}$ $\rightarrow r = \frac{2}{3}$ aef</p> <p>Substituting $\rightarrow a = \frac{3}{4}$ $\rightarrow S_{\infty} = \frac{\frac{3}{4}}{\frac{1}{3}} = 2\frac{1}{4}$ aef</p>	<p>M1</p> <p>A1</p> <p>M1 A1 [4]</p>	<p>Any valid method, seen or implied. Could be answers only.</p> <p>Both a and r</p> <p>Correct formula with $r < 1$, cao</p>
<p>(b)</p> <p>$4a = a + 4d \rightarrow 3a = 4d$</p> <p>$360 = S_5 = \frac{5}{2}(2a + 4d)$ or $12.5a$</p> <p>$\rightarrow a = 28.8^\circ$ aef Largest = $a + 4d$ or $4a = 115.2^\circ$ aef</p>	<p>B1</p> <p>M1</p> <p>A1 B1 [4]</p>	<p>May be implied in $360 = 5/2(a + 4a)$</p> <p>Correct S_n formula or sum of 5 terms</p> <p>cao, may be implied (may use degrees or radians)</p>

Page 7	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2015	9709	11

8	$f: x \mapsto 5 + 3\cos\left(\frac{1}{2}x\right)$ for $0 \leq x \leq 2\pi$.		
(i)	$5 + 3\cos\left(\frac{1}{2}x\right) = 7$ $\cos\left(\frac{1}{2}x\right) = \frac{2}{3}$ $\frac{1}{2}x = 0.84 \quad x = 1.68 \text{ only, aef}$ (in given range)	B1	Makes $\cos\left(\frac{1}{2}x\right) = \frac{2}{3}$
		M1A1 [3]	Looks up \cos^{-1} first, then $\times 2$
(ii)		B1	y always +ve, m always –ve. from (0, 8) to (2π, 2) (may be implied)
		B1	
		[2]	
(iii)	No turning point on graph or 1:1	B1	cao, independent of graph in (ii)
		[1]	
(iv)	$y = 5 + 3\cos\left(\frac{1}{2}x\right)$ Order; $-5, \div 3, \cos^{-1}, \times 2$ $x = 2\cos^{-1}\left(\frac{x-5}{3}\right)$	M1	Tries to make x subject.
		M1	Correct order of operations
		A1	cao
		[3]	

Page 8	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2015	9709	11

9	$y = x^3 + px^2$ (i) $\frac{dy}{dx} = 3x^2 + 2px$ Sets to 0 $\rightarrow x = 0$ or $-\frac{2p}{3}$ $\rightarrow (0, 0)$ or $\left(-\frac{2p}{3}, \frac{4p^3}{27}\right)$	B1 M1 A1 A1 [4]	cao Sets differential to 0 cao cao, first A1 for any correct turning point or any correct pair of x values. 2nd A1 for 2 complete TPs
	(ii) $\frac{d^2y}{dx^2} = 6x + 2p$ At $(0, 0) \rightarrow 2p$ +ve Minimum At $\left(-\frac{2p}{3}, \frac{4p^3}{27}\right) \rightarrow -2p$ -ve Maximum	M1 A1 A1 [3]	Other methods include; clear demonstration of sign change of gradient, clear reference to the shape of the curve www
(iii)	$y = x^3 + px^2 + px \rightarrow 3x^2 + 2px + p (= 0)$ Uses $b^2 - 4ac$ $\rightarrow 4p^2 - 12p < 0$ $\rightarrow 0 < p < 3$ aef	B1 M1 A1 [3]	Any correct use of discriminant cao (condone \leq)

Page 9	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2015	9709	11

<p>10</p> <p>(i)</p> <p>(ii)</p>	$y = \frac{8}{\sqrt{3x+4}}$ $\frac{dy}{dx} = \frac{-4}{(3x+4)^{\frac{3}{2}}} \times 3 \text{ aef}$ $\rightarrow m_{(x=0)} = -\frac{3}{2} \text{ Perpendicular } m_{(x=0)} = \frac{2}{3}$ <p>Eqn of normal $y - 4 = \frac{2}{3}(x - 0)$</p> <p>Meets $x = 4$ at $B \left(4, \frac{20}{3}\right)$</p> $\int \frac{8}{\sqrt{3x+4}} dx = \frac{8\sqrt{3x+4}}{\frac{1}{2}} \div 3$ <p>Limits from 0 to 4 \rightarrow Area $P = \frac{32}{3}$</p> <p>Area $Q =$ Trapezium $- P$ Area of Trapezium = $\frac{1}{2} \left(4 + \frac{20}{3}\right) \times 4 = \frac{64}{3}$</p> $\rightarrow \text{Areas of } P \text{ and } Q \text{ are both } \frac{32}{3}$	<p>B1 B1</p> <p>M1</p> <p>M1</p> <p>A1 [5]</p> <p>B1 B1</p> <p>M1 A1</p> <p>M1</p> <p>A1 [6]</p>	<p>Without the “×3” For “×3” even if 1st B mark lost.</p> <p>Use of $m_1 m_2 = -1$ after attempting to find $\frac{dy}{dx}_{(x=0)}$</p> <p>Unsimplified line equation</p> <p>cao</p> <p>Without “÷3”. For “÷3”</p> <p>Correct use of correct limits. cao</p> <p>Correct method for area of trapezium</p> <p>All correct.</p>
---	--	--	--