



Oxford Cambridge and RSA

GCE

Further Mathematics A

Y541/01: Pure Core 2

Advanced GCE

Mark Scheme for Autumn 2021

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2021

Annotations and abbreviations

Annotation in RM assessor	Meaning
✓ and ✖	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
^	Omission sign
MR	Misread
BP	Blank Page
NBOD	Benefit of doubt not given
Highlighting	
Other abbreviations in mark scheme	Meaning
dep*	Mark dependent on a previous mark, indicated by *. The * may be omitted if only one previous M mark
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working
AG	Answer given
awrt	Anything which rounds to
BC	By Calculator
DR	This question included the instruction: In this question you must show detailed reasoning.

Question		Answer	Marks	AO	Guidance	
1		$\begin{pmatrix} -12 & -1 & 5 \\ -1 & -20 & 3 \\ 7-6a & 3a-2 & -4a-5 \end{pmatrix}$ or $\begin{pmatrix} -4a & -1 & 5 \\ 11-4a & -20 & 3 \\ -8-a & 7 & -17 \end{pmatrix}$ seen $-12 = -4a$ or $-1 = 11 - 4a$ or $7 - 6a = -8 - a$ or $3a - 2 = 7$ or $-4a - 5 = -17$	M1	1.1	Either product AB or BA calculated (but not if assigned incorrectly). Alternatively: equivalent correct useful entries calculated for both	Condone 3 errors or omissions This mark can be implied by sight of a correct equation
			M1	1.1	Finding matrix products both ways and equating entries usefully	This mark can be implied by sight of a correct equation even if other entries or equations are wrong.
			A1	2.2a		Cannot be awarded if either AB or BA has more than 3 errors
		$a = 3$	[3]			

Question			Answer	Marks	AO	Guidance	
2	(a)	(i)	DR $3z_1 + 4z_2 = 3(3 - 7i) + 4(2 + 4i) = 17 - 5i$	B1 [1]	1.1		
		(ii)	DR $z_1z_2 = (3 - 7i)(2 + 4i) = 6 + 12i - 14i - 28(-1)$ $= 34 - 2i$	M1 A1 [2]	1.1 1.1	Attempted expansion with $i^2 = -1$ used and at least 3 correctly expanded terms	$-28(-1)$ can be simply +28
		(iii)	DR $\frac{z_1}{z_2} = \frac{3 - 7i}{2 + 4i} = \frac{3 - 7i}{2 + 4i} \times \frac{2 - 4i}{2 - 4i}$ $= \frac{6 - 12i - 14i - 28}{4 + 16} = \frac{-22 - 26i}{20} = -\frac{11}{10} - \frac{13}{10}i$	M1 A1 [2]	1.1 1.1	Multiplying top and bottom by (real multiple of) conjugate of bottom Must see some evidence of expansion	Allow $\frac{-11 - 13i}{10}$ or $-\frac{11 + 13i}{10}$
	(b)		DR $\sqrt{3^2 + (-7)^2}$ or $\tan^{-1}\left(\frac{-7}{3}\right)$ $ z_1 = \sqrt{58}$ or awrt 7.62 or $\arg z_1 = \text{awrt } -1.17$ or 5.12 rads $z_1 = \sqrt{58}\text{cis}(-1.17)$ or $z_1 = \sqrt{58}e^{-1.17i}$ or $z_1 = \sqrt{58}(\cos(-1.17) + i\sin(-1.17))$ or $[\sqrt{58}, -1.17]$	M1 A1 A1 [3]	1.1 1.1 2.5	Explicit working must be seen Must be in correct form with $\sqrt{58}$ exact and could be awrt 5.12 instead of -1.17 .	Other trig calculations could be sufficient for M1 provided that these are being used to find the argument. Do not condone degrees Condone round brackets

Question		Answer	Marks	AO	Guidance
3	(a)	$\left(\begin{pmatrix} 1 \\ -3 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 3 \\ 2 \\ -2 \end{pmatrix} \right) \cdot \begin{pmatrix} 2 \\ -5 \\ -3 \end{pmatrix} = 4$ $2 + 15 - 9 + \lambda(6 - 10 + 6) = 4$ $8 + 2\lambda = 4 \Rightarrow 2\lambda = -4 \Rightarrow \lambda = -2 \text{ so}$ $\mathbf{r} = \begin{pmatrix} 1 \\ -3 \\ 3 \end{pmatrix} + -2 \begin{pmatrix} 3 \\ 2 \\ -2 \end{pmatrix} = \begin{pmatrix} -5 \\ -7 \\ 7 \end{pmatrix}$	M1 M1 A1 [3]	1.1 1.1 1.1	Substituting the expression for a point on the line into the equation of the plane Dotting out to form and solve equation in λ Condone coordinates
	(b)	$\frac{\begin{pmatrix} 2 \\ -5 \\ -3 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 2 \\ -2 \end{pmatrix}}{\sqrt{4+25+9}\sqrt{9+4+4}} \text{ soi}$ $= \frac{6-10+6}{\sqrt{38}\sqrt{17}} = \frac{2}{\sqrt{646}} = 0.07868\dots$ $\theta = \text{awrt } 85.5\dots^\circ \text{ soi}$ $(\phi = 90^\circ - 85.48\dots^\circ =) \text{ awrt } 4.51^\circ$	M1 A1 A1 [3]	1.1 1.1 1.1	BC. Using $\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{ \mathbf{a} \mathbf{b} }$ Can be implied by correct final answer May see $\sin \phi = \frac{\mathbf{a} \cdot \mathbf{b}}{ \mathbf{a} \mathbf{b} }$ Or use of cross product or 1.49... rads or 0.0788 rads

Question		Answer	Marks	AO	Guidance	
3	(c)	$\lambda = 1 \Rightarrow \mathbf{r} = \begin{pmatrix} 4 \\ -1 \\ 1 \end{pmatrix}$ $\mathbf{b} = \begin{pmatrix} 3 \\ 2 \\ -2 \end{pmatrix} \times \begin{pmatrix} 2 \\ -5 \\ -3 \end{pmatrix} = \begin{pmatrix} -16 \\ 5 \\ -19 \end{pmatrix}$ <p>So equation of l_2 is $\mathbf{r} = \begin{pmatrix} 4 \\ -1 \\ 1 \end{pmatrix} + \mu \begin{pmatrix} -16 \\ 5 \\ -19 \end{pmatrix}$ oe</p>	B1 M1 A1 [3]	3.1a 2.2a 1.1	Method shown or at least two terms correctly evaluated Must be $\mathbf{r} =$. Allow parameter λ .	
4		DR $\sum_{r=1}^{100} (2r+3)^2 = 4 \sum_{r=1}^{100} r^2 + 12 \sum_{r=1}^{100} r + 9 \sum_{r=1}^{100} 1$ $\sum_{r=1}^{100} r^2 = \frac{1}{6} \times 100(100+1)(2 \times 100 + 1)$ $4 \times 338350 + 12 \times \frac{1}{2} \times 100 \times 101 + 900 = 1414900$	B1 M1 A1 [3]	3.1a 1.1a 1.1	Expanding and separating Use of formula for $\sum_{r=1}^{100} r^2$	

Question		Answer	Marks	AO	Guidance
5	(a)	DR $\text{RHS} = 2\cosh^2 x - 1 = 2\left(\frac{e^x + e^{-x}}{2}\right)^2 - 1$ $= 2\left(\frac{e^{2x} + 2 + e^{-2x}}{4}\right) - 1 = \frac{e^{2x} + 2 + e^{-2x}}{2}$ $= \frac{e^{2x} + e^{-2x}}{2} = \cosh 2x = \text{LHS}$	M1 A1 [2]	2.1 2.1	Uses correct exponential form in an attempt at proof AG Proof must be complete
	(b)	DR $2\cosh^2 x - 1 = 3\cosh x + 1$ $\Rightarrow 2\cosh^2 x - 3\cosh x - 2 = 0$ $(2\cosh x + 1)(\cosh x - 2) = 0$ $\cosh x = 2 \text{ or } -\frac{1}{2}$ $\cosh x \geq 1 \text{ so } \neq -\frac{1}{2}$ $x = \cosh^{-1} 2 = \ln(2 + \sqrt{3})$ $x = \ln(2 - \sqrt{3})$	M1 M1 A1 A1 A1 A1 [6]	3.1a 1.1 1.1 2.3 1.1 1.1	Use of identity in (a) to leave a three term quadratic equation in just cosh x Attempt to solve eg $\frac{-(-3) \pm \sqrt{(-3)^2 - 4 \times 2 \times (-2)}}{2 \times 2}$ Justification must be seen and must contain no incorrect statements For either correct answer seen Both correct values for x

		Answer	Marks	AO	Guidance	
6	(a)	DR A shear which leaves the x -axis invariant and which transforms the point $(0, 1)$ to the point $(2, 1)$.	B1 [1]	2.2a	Or any useful point transformed to its image	not “scale factor” or sf
	(b)	DR $\det A = 1 \times 1 - 0 \times 2 = 1$ and this is the area scale factor	B1 [1]	2.4	Both	Detailed calculation must be shown
	(c)	DR $\begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix}$ seen $\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix} = \begin{pmatrix} 7 & 2 \\ 3 & 1 \end{pmatrix}$ $\begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix} \begin{pmatrix} 7 & 2 \\ 3 & 1 \end{pmatrix}$ $= \begin{pmatrix} 7 & 2 \\ 3p & p \end{pmatrix} \Rightarrow p = 7$	B1 B1 M1 A1 [4]	3.1a 1.1 1.1 1.1	BC Correct form for stretch multiplied into their matrix in either order Correct multiplication	

Question		Answer	Marks	AO	Guidance
7	(a)	DR $\frac{x^3 + x^2 + 9x - 1}{x^3 + x^2 + 4x + 4} \equiv \frac{x^3 + x^2 + 4x + 4 + 5x - 5}{x^3 + x^2 + 4x + 4}$ $= 1 + \frac{5x - 5}{x^3 + x^2 + 4x + 4}$ So $A = 1, B = 5$ and $C = -5$	B1 [1]	3.1a	Attempt to divide out improper fraction. Could be by symbolic division or other valid method (eg comparing coefficients or substitution of values for x) Allow embedded answers
	(b)	DR $x^3 + x^2 + 4x + 4 = (x + 1)(x^2 + 4)$ $\frac{5x - 5}{x^3 + x^2 + 4x + 4} = \frac{D}{x + 1} + \frac{Ex + F}{x^2 + 4}$ $D(x^2 + 4) + (x + 1)(Ex + F) = 5x - 5$ $x = -1 \Rightarrow 5D = -10 \Rightarrow D = -2$ $x = 0 \Rightarrow -2 - F = -5 \Rightarrow F = 3$ $x^2 : D + E = 0 \Rightarrow E = 2$ $1 - \frac{2}{x + 1} + \frac{2x + 3}{x^2 + 4}$	B1 M1 A1 A1 A1 [5]	3.1a 1.2 1.1 1.1 1.1	Correct factorisation of cubic seen in working Correct form for partial fractions equated to their remainder rational fraction from (a). Follow through their division and factorisation. Or equivalent to find D correctly. Allow ft A1 for second and third coefficients found. Or $1 - \frac{2}{x + 1} + \frac{2x}{x^2 + 4} + \frac{3}{x^2 + 4}$

Question		Answer	Marks	AO	Guidance
7	(c)	DR $\int_0^2 \frac{x^3 + x^2 + 9x - 1}{x^3 + x^2 + 4x + 4} dx = \int_0^2 1 - \frac{2}{x+1} + \frac{2x}{x^2+4} + \frac{3}{x^2+4} dx$ $= \left[x - 2 \ln(x+1) + \ln(x^2+4) + \frac{3}{2} \tan^{-1}\left(\frac{x}{2}\right) \right]_0^2$ $\left(2 - 2 \ln 3 + \ln 8 + \frac{3\pi}{8} \right) - \ln 4$ $2 + \ln\left(\frac{2}{9}\right) + \frac{3}{8}\pi$	*M1 dep*M1 M1 A1 [4]	3.1a 1.1 1.1 1.1	Split term with $x^2 + 4$ in denominator and $ax + b$ in numerator Correctly integrate <i>their</i> expression (ignore limits) Correctly substitute limits to produce exact values and evaluate their \tan^{-1} term $a = 2, b = \frac{2}{9}, c = \frac{3}{8}$

8	(a)	$F = ma = 2 \frac{dv}{dt} = 4e^{-2t} - kv$	M1	3.3	Use of NII with m and a replaced and with 2 forces, the given force and kv	$F=ma$ can be implicit here
		$t = \ln 2, v = 0.5, F = 0 \Rightarrow 0 = 1 - 0.5k$	M1	2.2a	Use of given conditions to derive an equation in k	Can be done first
		$k = 2 \Rightarrow 2 \frac{dv}{dt} = 4e^{-2t} - 2v \Rightarrow \frac{dv}{dt} + v = 2e^{-2t}$	A1	1.1	AG	Complete argument including $F=ma$
		[3]				
	(b)	IF = $e^{\int 1 dt} = e^t$	*B1	1.1		Or CF
		$e^t \frac{dv}{dt} + e^t v = \frac{d}{dt}(e^t v) = e^t \times 2e^{-2t}$	*M1	1.1	Multiplying by IF and writing LHS as an exact derivative	Or subst correct PI into DE
		$e^t v = \int 2e^{-t} dt = -2e^{-t} + c$	A1	1.1	“+ c ” required	Or GS $v = Ae^{-t} - 2e^{-2t}$
		$t = 0, v = 0 \Rightarrow c = 2$	dep*M1	3.4	Use of initial conditions to derive a value for c	Or using alternative boundary condition
		$v = 2e^{-t} - 2e^{-2t}$	A1	3.4		
[5]						
	(c)	As $t \rightarrow \infty, v \rightarrow 0$	M1	3.4		
		So speed starts at 0 and ends at 0 (and is continuous and positive between) so must reach a maximum somewhere in $t > 0$	A1	2.4		
[2]						
	(d)	v is max when $\frac{dv}{dt} = 0$ so $t = \ln 2$	M1	2.2a	Deducing time when v is maximum	Or by finding expression for $\frac{dv}{dt}$
		So $v_{\max} = 0.5$ (given) (or	A1	3.4		and solving $\frac{dv}{dt} = 0$
		$v_{\max} = 2e^{-\ln 2} - 2e^{-2\ln 2} = 1 - \frac{2}{4} = \frac{1}{2}$)	[2]			

Question		Answer	Marks	AO	Guidance	
8	(e)	$v = \frac{dx}{dt} = 2e^{-t} - 2e^{-2t} \Rightarrow x = -2e^{-t} + e^{-2t} + d$ $t = 0, x = 0 \Rightarrow 0 = -2 + 1 + d \Rightarrow d = 1$ $0.9 = -2e^{-t} + e^{-2t} + 1$ $(e^{-t})^2 - 2e^{-t} + 0.1 = 0 \Rightarrow e^{-t} = \frac{10 \pm 3\sqrt{10}}{10}$ $\Rightarrow t = \ln\left(\frac{10}{10 - 3\sqrt{10}}\right) = 2.97 \text{ (3 sf)}$	M1 M1 M1 A1	3.3 3.3 3.5a 2.3	Integrating to find expression for x Using initial conditions to find value of (new) constant Recognising that the model is only valid when x lies between 0 and 0.9 Rejecting $t = \ln\left(\frac{10}{10 + 3\sqrt{10}}\right) < 0$ (can be implicit)	Or definite integral with correct lower limit.. ...and upper limit
9	(a)	$\mathbf{A}^2 = \begin{pmatrix} 4 & 12 \\ 0 & 4 \end{pmatrix}, \mathbf{A}^3 = \begin{pmatrix} 8 & 36 \\ 0 & 8 \end{pmatrix},$ $\mathbf{A}^4 = \begin{pmatrix} 16 & 96 \\ 0 & 16 \end{pmatrix}$ <p>Conjecture: $\mathbf{A}^n = \begin{pmatrix} 2^n & 3n \times 2^{n-1} \\ 0 & 2^n \end{pmatrix}$</p>	 B1 B1	 2.2a 2.2b	BC Allow this mark for any conjecture which works for $n = 1, 2, 3$ and 4.	

Question	Answer	Marks	AO	Guidance	
9	<p>(b)</p> <p>Basis case: $n = 1$:</p> $\mathbf{A}^1 = \begin{pmatrix} 2^1 & 3 \times 1 \times 2^0 \\ 0 & 2^1 \end{pmatrix} = \begin{pmatrix} 2 & 3 \\ 0 & 2 \end{pmatrix} = \mathbf{A} \text{ so true}$ <p>for $n = 1$</p> <p>Assume true for $n = k$</p> <p>ie $\mathbf{A}^k = \begin{pmatrix} 2^k & 3k \times 2^{k-1} \\ 0 & 2^k \end{pmatrix}$</p> $\mathbf{A}^{k+1} = \mathbf{A}^k \mathbf{A} = \begin{pmatrix} 2^k & 3k \times 2^{k-1} \\ 0 & 2^k \end{pmatrix} \begin{pmatrix} 2 & 3 \\ 0 & 2 \end{pmatrix}$ $= \begin{pmatrix} 2^{k+1} & 3 \times 2^k + 3k \times 2^k \\ 0 & 2^{k+1} \end{pmatrix}$ $= \begin{pmatrix} 2^{k+1} & 3(k+1) \times 2^k \\ 0 & 2^{k+1} \end{pmatrix}$ <p>So true for $n = k \Rightarrow$ true for $n = k + 1$. But true for $n = 1$.</p> <p>So true for all positive integer n</p>	<p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>[4]</p>	<p>2.1</p> <p>2.1</p> <p>2.2a</p> <p>2.4</p>	<p>Allow this mark even if the conjecture is wrong, provided that it works for $n = 1$</p> <p>Must have statement in terms of some other variable than n. Conjecture need not be correct.</p> <p>Uses inductive hypothesis properly & expands</p> <p>AG. Manipulating terms correctly and convincingly to obtain required form. Some intermediate working must be seen and a clear conclusion must be given for the induction process.</p>	<p>A formal proof by induction is required for full marks.</p>

Question		Answer	Marks	AO	Guidance
10	(a)	DR $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots = (1+x) + \left(\frac{x^2}{2!} + \frac{x^3}{3!} + \dots \right)$ $x > 0 \Rightarrow \frac{x^2}{2!} + \frac{x^3}{3!} + \dots > 0$ $\Rightarrow e^x > 1+x$	M1	1.1	Quoting and <i>using</i> the Maclaurin series
			A1	2.2a	AG. Result with sufficient justification
	(b)	DR $t = x+1 \Rightarrow e^{t-1} > t \Rightarrow \frac{e^t}{e} > t \Rightarrow e^t > et$	B1	3.1a	AG
	(c)	DR $t = \frac{\pi}{e} > 1 \text{ since } 2 < e < 3 \text{ and } \pi > 3$ $e^{\frac{\pi}{e}} > e \times \frac{\pi}{e} (= \pi)$ $\Rightarrow e^\pi > \pi^e \text{ (ie } e^\pi \text{ is greater)}$	B1	3.1a	Some justification that $t > 1$ is required
		M1	3.1a	Substituting their choice into the inequality	
		A1	1.1	Answer without use of inequality in part (b) scores M0A0	
		Alternative method $t = \ln \pi$ $e^{\ln \pi} > e \ln \pi$ $e^{\ln \pi} > e \ln \pi$ $\pi > \ln(\pi^e)$ $e^\pi > \pi^e$	B1		Some justification that $t > 1$ is required
			M1		
			A1		
			[3]		

OCR (Oxford Cambridge and RSA Examinations)
The Triangle Building
Shaftesbury Road
Cambridge
CB2 8EA

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored