



Rewarding Learning

**ADVANCED
General Certificate of Education
Reserve Summer 2022**

Mathematics

Assessment Unit A2 1

assessing

Pure Mathematics

[AMT11]

MONDAY 27 JUNE, MORNING

**MARK
SCHEME**

GCE ADVANCED/ADVANCED SUBSIDIARY (AS) MATHEMATICS

Introduction

The mark scheme normally provides the most popular solution to each question. Other solutions given by candidates are evaluated and credit given as appropriate; these alternative methods are not usually illustrated in the published mark scheme.

The marks awarded for each question are shown in the right-hand column and they are prefixed by the letters **M**, **W** and **MW** as appropriate. The key to the mark scheme is given below:

M indicates marks for correct method.

W indicates marks for working.

MW indicates marks for combined method and working.

The solution to a question gains marks for correct method and marks for an accurate working based on this method. Where the method is not correct no marks can be given.

A later part of a question may require a candidate to use an answer obtained from an earlier part of the same question. A candidate who gets the wrong answer to the earlier part and goes on to the later part is naturally unaware that the wrong data is being used and is actually undertaking the solution of a parallel problem from the point at which the error occurred. If such a candidate continues to apply correct method, then the candidate's individual working must be followed through from the error. If no further errors are made, then the candidate is penalised only for the initial error. Solutions containing two or more working or transcription errors are treated in the same way. This process is usually referred to as "follow-through marking" and allows a candidate to gain credit for that part of a solution which follows a working or transcription error.

Positive marking:

It is our intention to reward candidates for any demonstration of relevant knowledge, skills or understanding. For this reason we adopt a policy of **following through** their answers, that is, having penalised a candidate for an error, we mark the succeeding parts of the question using the candidate's value or answers and award marks accordingly.

Some common examples of this occur in the following cases:

- (a) a numerical error in one entry in a table of values might lead to several answers being incorrect, but these might not be essentially separate errors;
- (b) readings taken from candidates' inaccurate graphs may not agree with the answers expected but might be consistent with the graphs drawn.

When the candidate misreads a question in such a way as to make the question easier only a proportion of the marks will be available (based on the professional judgement of the examining team).

COVID-19 Context:

Given the unprecedented circumstances presented by the COVID-19 public health crisis, senior examiners, under the instruction of CCEA awarding organisation, are required to train assistant examiners to apply the mark scheme in case of disrupted learning and lost teaching time. The interpretation and intended application of the mark scheme for this examination series will be communicated through the standardising meeting by the Chief or Principal Examiner and will be monitored through the supervision period. This paragraph will apply to examination series in 2021-2022 only.

		AVAILABLE MARKS
1 (a) (i)	Diverges	MW1
	(ii) Oscillates	MW1
	(iii) Converges	MW1
(b)	$\sum_{r=1}^{\infty} 3 \left(\frac{2}{5}\right)^r = 3 \times \left(\frac{2}{5}\right) + 3 \times \left(\frac{2}{5}\right)^2 + 3 \times \left(\frac{2}{5}\right)^3 + \dots$ <p>This is a GP with $a = \frac{6}{5}$ $r = \frac{2}{5}$</p> $S_{\infty} = \frac{a}{1-r}$ $= \frac{\frac{6}{5}}{1 - \frac{2}{5}}$ $= 2$	M1 W2
(c) (i)	$3x^2 - (3x + 1) = 3x + 1 - x$ $3x^2 - 3x - 1 = 2x + 1$ $3x^2 - 5x - 2 = 0$ $(3x + 1)(x - 2) = 0$ $x = -\frac{1}{3}, x = 2$ <p>Hence $x = 2$</p>	M1 W1 MW1
	(ii) $d = 3x + 1 - x = 2x + 1$ $\Rightarrow d = 5$	W1 MW1
	$S_n = \frac{1}{2} n \{2a + (n - 1)d\}$ $= \frac{1}{2} \times 10 \{2(2) + 9(5)\}$ $= 245$	M1 W1 W1

16

		AVAILABLE MARKS
2	(i) $P = 2\pi r - r\left(\frac{\pi}{3}\right) + r$ $= \frac{5\pi r}{3} + r$	MW3
	(ii) $10\pi + 6 = r\left(\frac{5\pi}{3} + 1\right)$ $r = 6$	M1 W1
	(iii) Area of $\triangle AOB = \frac{1}{2} \times 6^2 \times \sin\left(\frac{\pi}{3}\right)$ $= 9\sqrt{3}$	MW1
	Area of minor sector AOB $= \frac{1}{2} \times 6^2 \times \frac{\pi}{3}$ $= 6\pi$	MW1
	Area of minor segment $= 6\pi - 9\sqrt{3}$	MW1
	Area of major segment $= \pi \times 6^2 - (6\pi - 9\sqrt{3})$ $= (30\pi + 9\sqrt{3}) \text{ cm}^2$	M1 W1
	3	
	(a) $\frac{4x^2 - 25}{3x^2 + 14x + 8} \times \frac{x + 4}{6x - 15}$ $\equiv \frac{(2x - 5)(2x + 5)}{(3x + 2)(x + 4)} \times \frac{x + 4}{3(2x - 5)}$ $\equiv \frac{2x + 5}{3(3x + 2)}$	M1 M1 W2 MW1
	(b) $(x + 3)(x + 2)^{-\frac{1}{2}}$ $= 2^{-\frac{1}{2}}(x + 3)\left(1 + \frac{x}{2}\right)^{-\frac{1}{2}}$ $= 2^{-\frac{1}{2}}(x + 3)\left(1 + \left(-\frac{1}{2}\right)\left(\frac{x}{2}\right) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2!}\left(\frac{x}{2}\right)^2 + \dots\right)$ $= \frac{1}{\sqrt{2}}(x + 3)\left(1 - \frac{x}{4} + \frac{3x^2}{32} + \dots\right)$ $= \frac{1}{\sqrt{2}}\left(x - \frac{x^2}{4} + 3 - \frac{3x}{4} + \frac{9x^2}{32} + \dots\right)$ $= \frac{1}{\sqrt{2}}\left(3 + \frac{x}{4} + \frac{x^2}{32} + \dots\right)$	M1 W1 M1 W1 M1 W1
		10
	11	

		AVAILABLE MARKS
4	(a) (i) Graph D	MW1
	(ii) Graph A	MW1
	(iii) Graph E	MW1
	(iv) Graph B	MW1
5	(b) (i) $1 \leq g(x) \leq 3, g(x) \in \mathbb{R}$	W2
	(ii) Let $y = \frac{1}{1+x}$	M1
	$\Rightarrow y(1+x) = 1$	M1
	$y + xy = 1$	
	$xy = 1 - y$	
	$x = \frac{1-y}{y}$	W1
	$\Rightarrow h^{-1}(x) = \frac{1-x}{x} \quad x \in \mathbb{R}, x \neq 0$	MW2
	(iii) $hg(x) = h(2 + \cos x)$	M1 W1
	$= \frac{1}{1+2+\cos x}$	W1
	$hg(x) = \frac{1}{3+\cos x} \quad x \in \mathbb{R}, \quad 0 \leq x \leq \pi$	MW1
(i) Let $f(x) = \operatorname{cosec}^2 3x - x^2 - 1$		
$f(0.3) = 0.53972\dots$	MW1	
$f(0.5) = -0.24497\dots$	MW1	
Since there is a change of sign in the value of $f(x)$ between $x = 0.3$ and $x = 0.5$, and $f(x)$ is continuous in this range, then there is a root between $x = 0.3$ and $x = 0.5$	MW1	
(ii) $f(x) = \operatorname{cosec}^2 3x - x^2 - 1$		
$f'(x) = 2 \operatorname{cosec} 3x \times (-3 \operatorname{cosec} 3x \cot 3x) - 2x$	MW1 W1	
$= -6 \operatorname{cosec}^2 3x \cot 3x - 2x$	W1	
$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$		
$x_1 = 0.3 - \left(\frac{\operatorname{cosec}^2(0.9) - 0.3^2 - 1}{-6 \operatorname{cosec}^2(0.9) \cot(0.9) - 0.6} \right)$	M1 W1	
$= 0.365 \text{ (3sf)}$	W1	

15

9

		AVAILABLE MARKS
6 (a)	$\text{LHS} \equiv \frac{2 \cos^2 \theta - 1 - \cos \theta + 1}{2 \sin \theta \cos \theta - \sin \theta}$ $\equiv \frac{2 \cos^2 \theta - \cos \theta}{2 \sin \theta \cos \theta - \sin \theta}$ $\equiv \frac{\cos \theta (2 \cos \theta - 1)}{\sin \theta (2 \cos \theta - 1)}$ $\equiv \frac{\cos \theta}{\sin \theta}$ $\equiv \cot \theta \equiv \text{RHS}$	M1 W1 MW1 M1 W1 MW1
(b)	$\tan(\theta - 45^\circ) = 6 \tan \theta$ $\Rightarrow \frac{\tan \theta - \tan 45^\circ}{1 + \tan \theta \tan 45^\circ} = 6 \tan \theta$ <p>Let $t = \tan \theta$</p> $\Rightarrow \frac{t - 1}{1 + t} = 6t$ $\Rightarrow t - 1 = 6t + 6t^2$ $6t^2 + 5t + 1 = 0$ $(3t + 1)(2t + 1) = 0$ $t = -\frac{1}{2}, -\frac{1}{3}$ $\tan \theta = -\frac{1}{2}$ $\theta = 153^\circ, 333^\circ$ $\tan \theta = -\frac{1}{3}$ $\theta = 162^\circ, 342^\circ$	M1 W1 MW1 MW1 W1 MW1 MW1

13

			AVAILABLE MARKS
7	(a)	$x = \sin t + \cos t$	M1
		$x^2 = (\sin t + \cos t)^2$	W1
		$x^2 = \sin^2 t + 2 \sin t \cos t + \cos^2 t$	W1
		$x^2 = 1 + \sin 2t$	
		$y = 4 - 3 \sin 2t$	
		$3 \sin 2t = 4 - y$	MW1
		$3(x^2 - 1) = 4 - y$	M1 W1
		$3x^2 = 7 - y$	
		$y = 7 - 3x^2$	W1
	(b)	Differentiate to give	
		$3 + 5y^2 + 10xy \frac{dy}{dx} = 4(x + y) \left(1 + \frac{dy}{dx}\right)$	M3 W2
		At (1, 3)	
		$3 + 45 + 30 \frac{dy}{dx} = 16 \left(1 + \frac{dy}{dx}\right)$	M1
		$\frac{dy}{dx}(30 - 16) = 16 - 48$	
		$\frac{dy}{dx} = -\frac{16}{7}$	W1
		Using (1, 3)	
		$y - 3 = -\frac{16}{7}(x - 1)$	M1
		$7y - 21 = -16x + 16$	
		$16x + 7y - 37 = 0$	W1
			16

8 (a) $V = \pi \int_0^{\frac{\pi}{16}} 9 \tan^2 4x \, dx$ M1 W2

$$= 9\pi \int_0^{\frac{\pi}{16}} (\sec^2 4x - 1) \, dx$$
M1 W1

$$= 9\pi \left[\frac{1}{4} \tan 4x - x \right]_0^{\frac{\pi}{16}}$$
W2

$$= 9\pi \left[\left(\frac{1}{4} \tan \left(\frac{\pi}{4} \right) - \frac{\pi}{16} \right) - (0 - 0) \right]$$
M1

$$= 9\pi \left(\frac{1}{4} - \frac{\pi}{16} \right)$$

$$= \frac{9\pi}{16} (4 - \pi) \text{ cubic units}$$
W1

(b) $\int x e^{2x} \, dx$

$$u = x \quad \frac{dv}{dx} = e^{2x}$$
M1 W1

$$\frac{du}{dx} = 1 \quad v = \frac{1}{2} e^{2x}$$
MW2

$$\Rightarrow \int x e^{2x} \, dx = \frac{1}{2} x e^{2x} - \int \frac{1}{2} e^{2x} \, dx$$
MW2

$$\int x e^{2x} \, dx = \frac{1}{2} x e^{2x} - \frac{1}{4} e^{2x} + c$$
W1

(c) $\int_4^6 \frac{x(x-5)}{(x-3)^2} \, dx$

$$u = x - 3 \Rightarrow x = u + 3$$

$$\frac{du}{dx} = 1$$
MW1

$$x = 4 \quad u = 1$$
M1

$$x = 6 \quad u = 3$$
W1

$$\Rightarrow \int_1^3 \frac{(u+3)(u-2)}{u^2} \, du$$
M1 W1

$$= \int_1^3 \frac{u^2 + u - 6}{u^2} \, du$$

$$= \int_1^3 \left(1 + \frac{1}{u} - \frac{6}{u^2} \right) \, du$$
W1

$$= \left[u + \ln|u| + \frac{6}{u} \right]_1^3$$
MW1

$$= [3 + \ln 3 + 2] - [1 + \ln 1 + 6]$$
M1

$$= -2 + \ln 3$$
W1

9 (i) Let $f(x) = 2x^3 + 11x^2 + 12x - 9$
 $f(-3) = 2(-27) + 11(9) + 12(-3) - 9$
 $= 0$
 $\Rightarrow (x + 3)$ is a factor

M1 W1

MW1

(ii)
$$x + 3 \overline{) 2x^3 + 11x^2 + 12x - 9}$$

$$\underline{2x^3 + 6x^2}$$

$$5x^2 + 12x$$

$$\underline{5x^2 + 15x}$$

$$-3x - 9$$

$$\underline{-3x - 9}$$

$$0$$

M1

$\Rightarrow (x + 3)(2x^2 + 5x - 3)$ W1

$\Rightarrow (x + 3)(2x - 1)(x + 3)$

$\Rightarrow f(x) = (x + 3)^2(2x - 1)$ MW1

$\Rightarrow \int \frac{x^2 + 3x + 35}{2x^3 + 11x^2 + 12x - 9} dx = \int \frac{x^2 + 3x + 35}{(x + 3)^2(2x - 1)} dx$ MW1

$\frac{x^2 + 3x + 35}{(x + 3)^2(2x - 1)} \equiv \frac{A}{x + 3} + \frac{B}{(x + 3)^2} + \frac{C}{2x - 1}$ M1 W1

Comparing numerators

$\Rightarrow A(x + 3)(2x - 1) + B(2x - 1) + C(x + 3)^2 \equiv x^2 + 3x + 35$ M1

$x = \frac{1}{2} \quad \frac{49}{4}C = \frac{147}{4} \quad \Rightarrow C = 3$ M1 W1

$x = -3 \quad -7B = 35 \quad \Rightarrow B = -5$ W1

x^2 term $2A + C = 1 \quad \Rightarrow A = -1$ M1 W1

$\Rightarrow \int \left(\frac{-1}{x + 3} + \frac{-5}{(x + 3)^2} + \frac{3}{2x - 1} \right) dx$

$= -\ln|x + 3| + \frac{5}{x + 3} + \frac{3}{2} \ln|2x - 1| + c$ M1 W2

AVAILABLE
MARKS

18

		AVAILABLE MARKS
<p>10 (i) $\frac{dA}{dt} = 0.25A - 50$</p> <p>$\Rightarrow \int \frac{dA}{0.25A - 50} = \int dt$</p> <p>$\Rightarrow 4 \ln(0.25A - 50) = t + c$</p> <p>$\ln(0.25A - 50) = 0.25t + 0.25c$</p> <p>$\Rightarrow 0.25A - 50 = ke^{0.25t}$</p> <p>$\Rightarrow A = 4ke^{0.25t} + 200$</p>	<p>M1 W1</p> <p>M1 W1</p> <p>MW2</p> <p>W1</p>	
<p>(ii) $t = 0, A = 190$</p> <p>$\Rightarrow 190 = 4k + 200$</p> <p>$4k = -10$</p> <p>$A = 200 - 10e^{0.25t}$</p>	<p>M1</p> <p>W1</p>	
<p>(iii) $A = 0, t = ?$</p> <p>$200 - 10e^{0.25t} = 0$</p> <p>$e^{0.25t} = 20$</p> <p>$0.25t = \ln 20$</p> <p>$t = 11.98$</p> <p>Minimum length of time for shift is 12 hours.</p>	<p>M1</p> <p>W1</p> <p>W1</p>	
<p>(iv) $t = 0, A = A_0$</p> <p>$\Rightarrow A_0 = 4k + 200$</p> <p>$\Rightarrow A = (A_0 - 200)e^{0.25t} + 200$</p> <p>$t = 8, A \leq 0$</p> <p>$(A_0 - 200)e^2 + 200 \leq 0$</p> <p>$A_0 \leq \frac{200e^2 - 200}{e^2}$</p> <p>$A_0 \leq 172.9$</p> <p>Maximum amount is 172.9 kg (1dp).</p>	<p>M1 W1</p> <p>W1</p>	
Total		17
		150