



**GCE AS/A level**

0979/01

**MATHEMATICS – FP3**  
**Further Pure Mathematics**

A.M. TUESDAY, 24 June 2014

1 hour 30 minutes

#### **ADDITIONAL MATERIALS**

In addition to this examination paper, you will need:

- a 12 page answer book;
- a Formula Booklet;
- a calculator.

#### **INSTRUCTIONS TO CANDIDATES**

Use black ink or black ball-point pen.

Answer **all** questions.

Sufficient working must be shown to demonstrate the **mathematical** method employed.

#### **INFORMATION FOR CANDIDATES**

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

1. (a) Starting with the exponential definition of  $\sinh x$ , show that

$$\sinh^{-1} x = \ln(x + \sqrt{x^2 + 1}). \quad [4]$$

- (b) Solve the equation

$$\cosh 2x = 2\sinh x + 5,$$

giving your answers in the form  $\ln(a + \sqrt{b})$  where  $a, b$  are integers. [5]

2. The equation  $x^3 + x = 3$  has a root  $\alpha$  between 1.2 and 1.3.

- (a) Alun suggests the following iterative sequence for finding the value of  $\alpha$  based on rearranging the equation

$$x_{n+1} = \sqrt[3]{3 - x_n} \text{ with } x_0 = 1.25.$$

By evaluating an appropriate derivative, show that this sequence is convergent. Use it to find the value of  $\alpha$  correct to 4 decimal places. [8]

- (b) Starting with  $x_0 = 1.25$ , use the Newton-Raphson method to find the value of  $\alpha$  correct to 6 decimal places. [6]

3. (a) Assuming the derivative of  $\cosh x$ , show that

$$\frac{d}{dx}(\operatorname{sech} x) = -\operatorname{sech} x \tanh x. \quad [1]$$

- (b) Determine the Maclaurin series for  $\tanh x$  as far as the term in  $x^3$ . [6]

- (c) Hence find an approximate value for the integral

$$\int_0^{0.5} (1+x)\tanh x \, dx.$$

Give your answer correct to three significant figures. [4]

4. Using the substitution  $t = \tan\left(\frac{x}{2}\right)$ , determine the value of the integral

$$\int_0^{\frac{\pi}{2}} \frac{1}{2 - \cos x} \, dx. \quad [8]$$

5. The integral  $I_n$  is defined, for  $n \geq 0$ , by

$$I_n = \int_0^1 x^n e^{-x^2} dx.$$

- (a) Show that, for  $n \geq 2$ ,

$$I_n = \left(\frac{n-1}{2}\right)I_{n-2} - \frac{e^{-1}}{2}. \quad [3]$$

- (b) Evaluate  $I_5$ , giving your answer in the form  $a - be^{-1}$ , where  $a, b$  are positive constants to be determined. [6]

6. The curve  $C$  has polar equation

$$r = \sin\theta + \cos\theta, \quad 0 \leq \theta \leq \frac{\pi}{2}.$$

- (a) Find the polar coordinates of the point at which the tangent is parallel to the initial line. [8]
- (b) Find the area of the region enclosed between  $C$ , the initial line and the line  $\theta = \frac{\pi}{2}$ . [5]

7. (a) Using the substitution  $x = a \sinh\theta$ , show that

$$\int \sqrt{x^2 + a^2} dx = \frac{a^2}{2} \left( \sinh^{-1}\left(\frac{x}{a}\right) + \frac{x\sqrt{x^2 + a^2}}{a^2} \right) + \text{constant}. \quad [5]$$

- (b) The equation of the curve  $C$  is

$$y = x^2, \quad 0 \leq x \leq 1.$$

Find the arc length of  $C$ . [6]

**END OF PAPER**